

Simulating the germination response to diurnally alternating temperatures under climate change scenarios: comparative studies on *Carex diandra* seeds

Eduardo Fernández-Pascual¹, Charlotte E. Seal^{2*}, Hugh W. Pritchard²

¹ Jardín Botánico Atlántico, Universidad de Oviedo, Avda. del Jardín Botánico 2230, 33394 Gijón/Xixón, Spain; ² Seed Conservation Department, Royal Botanic Gardens, Kew, Wellcome Trust Millennium Building, Wakehurst Place, Ardingly, West Sussex RH17 6TN, UK*

For correspondence. Charlotte E. Seal; Fax: +44 (0)1444 894110 Email: c.seal@kew.org

This is a pre-copyedited, author-produced PDF of an article accepted for publication in *Annals of Botany* following peer review. The version of record (Fernández Pascual, E., Seal, C.E. & Pritchard, H.W. (2015) Simulating the germination response to diurnally alternating temperatures under climate change scenarios: comparative studies on *Carex diandra* seeds. *Annals of Botany* 115: 201-209) is available online at: <http://dx.doi.org/10.1093/aob/mcu234>.

ABSTRACT

Background and Aims Environmental temperature regulates plant regeneration by seed in several superimposed ways, and this complex regulation will be disrupted by climate change. The role of diurnally alternating temperatures (ΔT) in terminating dormancy will be a major player in this disruption, as its effects on seed germination are immediate.

Methods We modelled the effect of ΔT on seed germination comparing two populations of the wetland sedge *Carex diandra*. We fitted a cardinal-temperature model to germination results in a thermal gradient plate, and used the model to simulate changes in germination under two representative concentration pathways scenarios (RCP2.6 and RCP8.5).

Key Results Increasing ΔT decreased the base temperature for germination and the thermal time required for germination. RCP2.6 projected moderate increases in average temperatures and ΔT , whereas RCP8.5 projected higher warming and ΔT . The higher ΔT effect together with the warmer temperatures increased seed germination in both scenarios. By considering the effects of ΔT , the model projects considerable increases in germination.

Conclusions *Carex diandra* germination will be very responsive to potential changes in ΔT as a consequence of climate change. This research highlights the role of ΔT in seed responses to climate change. Comprehensive cardinal-temperature models, encompassing the different effects of temperature on seed germination, are needed to understand how climate change will affect plant regeneration.

Key words: Cardinal-temperature models, germination thermal thresholds models, physiological dormancy, plant-climate interactions, plant regeneration, representative concentration pathways, seed ecology

INTRODUCTION

Climate change (IPCC, 2013) will disrupt the many interactions between biology and climate, from enzymatic reactions to ecological patterns. Climate thoroughly controls key processes such as plant regeneration, as is exemplified by the thermal regulation of seed germination (Probert, 2000, Fenner and Thompson, 2005). Temperature drives local adaptation (Wagmann et al., 2012, Fernández-Pascual et al., 2013a) and phenotypic plasticity (Chiang et al., 2011, Hoyle et al., 2008, Figueroa et al., 2010, Fernández-Pascual and Jiménez-Alfaro, 2014, Fenner, 1991) in germination traits, as well as the physiological processes of dormancy loss (Finch-Savage and Leubner-Metzger, 2006, Batlla and Benech-Arnold, 2005, Batlla and Benech-Arnold, 2003, Batlla and Benech-Arnold, 2004, Steadman and Pritchard, 2004, Pritchard et al., 1996) and germination elicitation (Hardegree, 2006, Pritchard and Manger, 1990, García-Huidobro et al., 1982). In seasonal climates, germination traits interplay with annual temperature cycles to ensure that seed emergence and seedling

establishment occur in the most favourable season (Donohue et al., 2010, Vleeshouwers et al., 1995). Given the significance of germination in the life history of a plant, it is not surprising that its timing is a central scenario for natural selection (Donohue et al., 2005a, Donohue et al., 2005b, Donohue et al., 2005c). However, the complex thermal control of germination timing is very responsive to climate change (Walck et al., 2011). New environmental temperatures may not match the temperatures that alleviate dormancy (Orrù et al., 2012) and elicit germination (Cochrane et al., 2011). This mismatch could alter recruitment from the soil seed bank (Hoyle et al., 2013, Ooi et al., 2009, Ooi, 2012) and shift germination timing (Mondoni et al., 2012), compromising plant regeneration and community composition.

Nonetheless, many aspects of the potential seed response to climate change remain unstudied. Together with a steady increase in average temperatures, climate change projections point to more frequent hot and fewer cold daily extremes (IPCC, 2013). Depending on the relative change in daily maxima and minima, diurnal

thermal oscillations could actually rise, fall or remain unchanged; affecting the physiological processes of species (Vasseur et al., 2014). It has been known for a long time that diurnally alternating temperatures (hereafter ΔT) promote germination in many plant species (Morinaga, 1926, Probert, 2000, Fenner and Thompson, 2005). This germination response to ΔT is especially common in wetland and disturbed habitats, where it detects favourable establishment environments with regard to burial depth, soil perturbation or water level (Thompson and Grime, 1983, Liu et al., 2013). Physiologically, ΔT may act through a decrease in the proportion between abscisic acid and gibberellins, which reduces the embryo's water potential and promotes radicle growth (Huarte and Benech-Arnold, 2010, Huarte and Benech-Arnold, 2005, Huarte, 2013). The effect of thermal oscillations is almost immediate, and on the first day after transfer from constant to ΔT regimes imbibed seeds undergo a large transcriptome change involving metabolism, plant hormones and the circadian clock (Foley et al., 2010). ΔT is thus a 'immediate' signal for dormancy alleviation which marks that emergence conditions are immediately favourable, contrasting with slower dormancy loss associated to seasonal change, e.g. overwintering (Finch-Savage and Leubner-Metzger, 2006). This immediacy means that if climate change modifies ΔT , plant regeneration will be strongly affected as small changes in amplitude will translate into quick changes in emergence.

Recent research has begun to explore seed responses to climate change using cardinal-temperature models (Orrù et al., 2012, Porceddu et al., 2013). In this approach, germination experiments are employed to fit germination models based on two thresholds, (1) the base temperature (T_b) below which the germination rate is zero and (2) the thermal time (θ) or degrees day above the T_b required to reach successive percentage germination increases in a seed population (Porceddu et al., 2013, Hardegree, 2006). Subsequently the models are tested under average temperature projections to simulate potential emergence patterns in the future (Orrù et al., 2012). In terms of cardinal-temperature models, increasing ΔT decreases the base temperature and the thermal time for germination, i.e. germination is faster (Qiu et al., 2006), similarly to other dormancy breaking factors such as chilling (Steadman and Pritchard, 2004, Pritchard et al., 1996, Pritchard et al., 1999). It is therefore clear that in those species with a germination response to ΔT , an accurate simulation will require the incorporation of the dormancy breaking effect of ΔT into the models. To our knowledge this step in model design and testing has not yet been taken.

Here we describe for the first time a simulation of seed germination under projected diurnal thermal oscillations according to climate change scenarios. To do so we chose as a study species the wetland sedge *Carex diandra* Schrank (Cyperaceae). *C. diandra* is a widely-distributed northern hemisphere sedge that occupies

transitional mire soils (Habitats Directive 7140 Transition mires and quaking bogs) in the margins of lakes and ponds (Castroviejo et al., 2007, Tutin et al., 1980), habitats where the germination response to ΔT seems to be a pervasive adaptation. We used seeds from two contrasting altitudes to establish comparative cardinal-temperature models describing the fresh germination response to ΔT . Then we used the models to simulate germination changes under the ΔT projected in representative concentration pathways (RCP) scenarios (Vuuren et al., 2011). RCPs define different pathways of total radiative forcing produced by accumulation of all sources of human greenhouse gas emissions. RCPs do not represent specific scenarios of environmental/economic projections; instead each of them could result from different combinations of economic, technological, demographic, policy, and institutional futures (http://sedac.ipcc-data.org/ddc/ar5_scenario_process/RCPs.html). We chose RCP2.6 (lowest radiative forcing, i.e. lowest climatic alteration) and RCP8.5 (highest radiative forcing, i.e. highest climatic alteration). We address two specific hypotheses in freshly dispersed seeds with conditional dormancy: (1) increasing ΔT will lower the base temperature and shorten the thermal time for germination; and (2) simulated germination timing will change under the ΔT projections of the RCPs.

MATERIALS AND METHODS

Study system

Carex diandra produces abundant seeds that disperse in summer and show conditional physiological dormancy, being able to germinate at warm temperatures immediately after dispersal and acquiring the capacity to germinate at cooler temperatures as they lose dormancy (Fernández-Pascual et al., 2013b). In late July 2012, ripe *C. diandra* seeds were collected from two ponds of the Cantabrian Mountains, the montane site 'La Mina' (43° 16' N 4° 59' W, 1110 m asl) and the subalpine site 'La Recoleta' (43° 1' N 6° 7' W, 1780 m asl) (Molina et al., 2009). The day after collection, seeds were cleaned and kept at controlled conditions (15 °C, 15 % relative humidity (RH)) until further use.

To calculate present and future values of environmental average temperatures and ΔT at the population sites, present model-derived monthly maximum and minimum temperatures at the population sites were downloaded from WorldClim (Hijmans et al., 2005); and projections of the same variables for the year 2070 under the RCP2.6 and RCP8.5 scenarios were downloaded from the HadGEM2-ES model (Jones et al., 2011).

Germination experiment

The germination experiment began in November 2012. For each population, 25 seeds were sown onto 169 Petri dishes (1 % agar). Each population was placed on a bidirectional thermal gradient plate (Model GRD1, Grant Instruments, Cambridge, UK). While one

gradient remained constant, the other changed direction every 12 h, providing 169 thermoperiod combinations with averages from 7.5 to 37.5 °C and ΔT from 0 to 30 °C (Fig. 1). The thermoperiods were coupled with a 12 / 12 light-darkness photoperiod, half of the dishes having a warmer day regime and half a warmer night regime. Actual temperatures in the gradient were recorded with five temperature probes (one in the centre and one in each corner of the thermal gradient plate), placed into the agar Petri dishes and set to record every 10 minutes for the duration of the experiment. From these records we calculated the average day and night temperatures of each Petri dish, and used these averages in further calculations. Germination (radicle emergence) was scored daily, and the dishes replenished periodically to prevent the agar from drying. The experiment lasted 5 weeks by which time no further germination was observed. A random sample ($n = 300$ per population) of non-germinated seeds were cut open to determine the occurrence of empty and fungus infected seeds. The percentage on non-viable seeds was low (4 %) and therefore the germination percentages were calculated over the full number of seeds sown (25).

Cardinal-temperature model

Cardinal-temperature models (García-Huidobro et al., 1982, Orrù et al., 2012, Hardegree, 2006, Pritchard and Manger, 1990, Porceddu et al., 2013, Qiu et al., 2006) were used to describe the germination thermal thresholds under different ΔT regimes. For each dish, the time (tg, days) taken for cumulative germination to reach successive 10 % increases (g) was calculated by linear interpolation on the germination curves. Data from dishes were grouped according to their regime of ΔT (± 2 °C), and for each regime the rate of germination ($1/tg$) was plotted against the average germination temperature. Available treatments did not allow analysis of the supra-optimal temperature range, which in any case is not relevant ecologically for the environment of the Cantabrian Mountains (see discussion). The suboptimal germination temperature range was identified by visual inspection of the germination rate plots, and a linear regression was fitted to it according to the equation:

$$1 / tg (d-1) = (T - T_b) / \theta g$$

The average x-intercept among g percentiles was the T_b for each ΔT , and T_b was allowed to vary among g to consider variation among individual seeds.

Subsequently the thermal time required to reach successive g (θg , °Cd) was estimated as the inverse of the slope of the sub-optimal linear regression equations. Values of g were transformed into probits, and new linear regressions were fitted between probit (g) and θg expressed as both linear θg and $\log \theta g$. Linear θg gave higher R^2 values, and therefore linear equations were used to describe the germination of seeds as a function of thermal time under each ΔT regime:

$$\text{probit (g)} = K + \theta g / \sigma$$

The reciprocal of the slope (σ) represents the sensitivity of the population to thermal time; flatter slopes indicating higher variability among individual seeds in their response. K is the intercept when θ is zero.

Statistical analysis

To determine how the thermal thresholds varied depending on the ΔT regime, the seed population and the phase when temperatures were warmer (day or night) Generalized Linear Models (gamma regression) were fitted to the values of T_b and $\theta 50$. Then, considering only the dishes with a warmer day regime (the ecologically meaningful situation) and separately for each population, linear or logarithmic equations were used to estimate T_b and $\theta 50$ as a function of ΔT values at present and in the RCP2.6 and RCP8.5 scenarios. Finally, the thermal thresholds and the accumulation of degrees day for germination were simulated under (1) present average monthly temperatures and ΔT ; (2) RCP average monthly temperatures but present ΔT ; (3) RCP average monthly temperatures and ΔT .

RESULTS

Germination response to temperature

Germination of *C. diandra* did not occur under constant thermoperiods, a ΔT of at least 2 °C was necessary for some germination to take place (Fig. 1). The highest germination percentages occurred with an average (day + night) temperature of around 25 °C and a ΔT of 10-14 °C. The montane population had higher germination (a total of 1798 germinated seeds) than the subalpine (687 germinated seeds). These differences were due to dormancy and not seed quality, as the subalpine seeds were able to produce >95 % germination in some treatments (e.g. 35/20 °C). Germination was possible both under warmer day and warmer night regimes, and although slightly more seeds germinated when the warm phase coincided with light, the differences were minimal (1258 seeds vs. 1227 seeds, averaging both populations).

Cardinal-temperature model and effect of ΔT

The T_b was significantly different among the two populations, being estimated at 16.8 °C ± 0.5 for the montane seeds and 20.3 °C ± 0.6 for the subalpine; whereas the phase of warmer temperatures did not affect the T_b significantly (Table 1). Conversely, the $\theta 50$ was not significantly different among populations but was significantly higher in warmer day regimes (258.8 °Cd ± 27.5), compared to warmer night regimes (162.1 °Cd ± 19.0) (Table 1).

We were able to establish sub-optimal cardinal-temperature models for ΔT regimes from 2 to 24 °C. ΔT significantly influenced both T_b and $\theta 50$ (Table 1). In both cases the effect followed a logarithmic function (Fig. 2), with a sharp decrease as ΔT increased from 0 to 10-14 °C and stabilization with higher values. The T_b followed a linear decrease between 0 and 14 °C; as this

matched the range of ΔT found in the temperatures of the models, we used a linear equation to estimate the monthly T_b under the monthly ΔT projected at present and under the two RCPs. For the $\theta 50$, we estimated an annual average using a logarithmic equation and the annual average ΔT under each scenario.

Simulated germination under climate change

The average annual temperature at present according to models was 6.7 °C with a ΔT of 9.5 °C. The RCP2.6 projected an average of 8.8 °C with a ΔT of 9.9 °C. The RCP8.5 projected an average of 11.0 °C with a ΔT of 10.3 °C. As a consequence of the increasing ΔT of the RCPs, T_b (averaging the two populations) was projected to decrease from 18.6 °C at present to 18.2 °C in RCP2.6 and to 17.8 °C in RCP 8.5 (Fig. 3). Present T_b and environmental temperatures would allow some accumulation of °C d for fresh germination at the montane site, but prevent it at the subalpine (Fig. 4). Under RCP2.6, the accumulation of °C d would increase notably in the montane population, but would still not occur in the subalpine (Fig. 4). Under RCP8.5 the accumulation would be high in both populations (Fig. 4). When the lowering of the T_b as a consequence of the new ΔT is incorporated into the simulation, the accumulation of °C d increases in both sites and RCPs, the effect being more marked in the subalpine site where accumulation appears in the RCP2.6 and doubles in the RCP8.5 (Fig. 4).

The $\theta 50$ would also vary with the ΔT of each climatic scenario, although the change would be minimal (Fig. 4). At the montane site it was projected to decrease by 4 % from the present value in the RCP2.6 and by 8 % in the RCP8.5. At the subalpine site the decrease would be comparatively smaller, 3% in the RCP2.6 and in the 5 % RCP8.5. Nonetheless, the combined effects of a higher accumulation of °C d and a lower $\theta 50$ would change seed germination patterns in both sites (Fig. 4). In the montane site, 50 % fresh germination would occur in the summer season of dispersal under either scenario. The subalpine population would be unaffected under the RCP 2.6, but under the RCP8.5 significant fresh germination would occur at dispersal, 50 % if the ΔT effect is considered.

DISCUSSION

Germination response to ΔT

Germination in *C. diandra* has an absolute requirement for ΔT . Such a strong germination response to ΔT is a common trait of sedges (Schütz, 2000, Schütz, 1997, Schütz and Rave, 1999, Budelsky and Galatowitsch, 1999) and wetland species in general (Thompson and Grime, 1983, Baskin and Baskin, 1998, Patzelt et al., 2001, Maas, 1989, Carta et al., 2013). In the case of *C. diandra*, it could allow seeds to detect the fall of the water table in the pond margins it colonizes. ΔT marks the beginning of the favourable season, at least in related fen soils of the Cantabrian Mountains where amplitude is minimal in winter, increases in spring and

peaks in summer (Fernández-Pascual et al., 2014). Nevertheless, there are differences among sites and ΔT is less successful in alleviating dormancy in the subalpine population, following a pattern of deeper dormancy at higher altitudes that is common to many species (Cavieres and Arroyo, 2000, Fernández-Pascual et al., 2013a, Zhou and Bao, 2014). Conditional dormancy in *C. diandra* can also be alleviated in the long term by cold-overwintering (Fernández-Pascual et al., 2013b), and this signal of slow seasonal change (Finch-Savage and Leubner-Metzger, 2006) is probably more significant in the subalpine site, where winter is harsher and premature emergence riskier.

These dormancy differences operate through T_b , mirroring the positive latitudinal correlation that has been found in the T_b of *Aesculus hippocastanum* L. populations (Daws et al., 2004). On the other hand, $\theta 50$ does not vary among populations, contrasting with the negative correlation between altitude and $\theta 50$ that has been found in *Vitis vinifera* L. (Orrù et al., 2012). The T_b , as well as the total number of germinated seeds, does not vary depending on whether the light phase coincides with the warm or the cool phase of the thermoperiod. This agrees with previous work that shows that the amplitude of the thermal oscillation is the main parameter that explains the germination response to ΔT (Roberts and Totterdell, 1981, Benech-Arnold et al., 1990a, Benech-Arnold et al., 1990b, Batlla et al., 2003). Conversely, the $\theta 50$ is significantly lower when the night is warmer than the day, i.e. seeds germinate faster in that case. It is difficult to interpret this abnormal and ecologically unrealistic situation, but it could be related to the role of the phytochrome in the response to ΔT (Probert, 2000).

A main effect of ΔT in *C. diandra* is that as ΔT increases, T_b decreases, thus facilitating the accumulation of °C d for germination (i.e., under a given average temperature, more degrees above the T_b can be accumulated each day). The relationship between T_b and ΔT is linear up to 10-14 °C, whereas it becomes asymptotic with larger ΔT values, as was found in certified seeds of *Dactylis glomerata* L. (Qiu et al., 2006). ΔT above the optimal amplitude did have a negative effect in wild European populations of *D. glomerata* (Probert et al., 1986), but in any case supraoptimal values do not seem realistic in *C. diandra* field conditions and therefore lack ecological significance; temperatures over 25 °C will not be reached in the Cantabrian Mountains even under the more extreme climatic scenarios (Álvarez García et al., 2009). The optimal ΔT in *C. diandra* and *D. glomerata* contrasts with what was found in grassland species living in disturbed soils at very high altitudes in the Tibetan Plateau, where the optimum ΔT was 20 °C and 10 °C had no effect; but such high amplitudes can be expected in the extreme Tibetan conditions (Liu et al., 2013). Therefore, the optimal ΔT might be closely related to the thermal range of the specific habitat.

Implications for climate change

The simulation of *C. diandra* germination under RCP scenarios points to an increase in emergence parallel to the intensity of the RCP scenario and to the associated increase in average temperatures. However, the full magnitude of the change is only appreciated when the response to ΔT is incorporated into the simulation, as this increases the daily accumulation of °C d at the same time that it decreases the cumulative °C d needed for the population to germinate. The effect of incorporating ΔT is especially relevant at the subalpine site, where the accumulation of °C d doubles under the RCP8.5 scenario. Total germination increases are higher at the montane site, where a high percentage of the seeds are not expected to overwinter and join the soil seed bank even under the moderate RCP2.6 scenario. But as germination at dispersal at the montane site seems to be possible at present, it is to be expected that seedlings might survive winters there. Conversely, the harsher winters at the subalpine site make premature germination in autumn or during early winter a more deleterious trait, and accordingly dormancy there is deeper. Therefore, the change could have more negative consequences at the subalpine site in the RCP8.5 scenario, as germination in the same summer of dispersal would change from negligible to affecting half the seed population. Whilst our findings are based on a simulation of how one species might respond across two sites, they do highlight the probable importance of ΔT on regeneration under climate change of the many plant species that show the response to ΔT . The latest climate change analysis at the *C. diandra* study sites report a higher increase of daily maxima in comparison with daily minima (Álvarez García et al., 2009), so the ΔT could be modifying germination patterns already.

It must be remembered that our simulation is a simplification of nature that concentrates on the effect of ΔT on fresh germination, but in the field temperature will influence germination in other ways. A new climate will modify germination traits through phenotypic plasticity (Fenner, 1991, Hoyle et al., 2008, Chiang et al., 2011, Fernández-Pascual and Jiménez-Alfaro, 2014, Fernández-Pascual et al., 2013a, Daws et al., 2004). For example, present differences in T_b between the montane and subalpine sites are larger (16.8 vs. 20.3 °C) than the differences in T_b projected among scenarios (18.6 to 17.8 °C). Therefore, if all the present variation among sites is attributable to phenotypic plasticity, this plasticity would accommodate the effect produced by climate change (Cochrane et al., 2014). However, our experimental design does not at this stage allow for differentiation between phenotypic plasticity and genetically-based local adaptation (Fernández-Pascual et al., 2013a).

Furthermore, our simulation is only valid for the germination of fresh seeds, i.e. seed germination between dispersal in summer and the beginning of the winter season in November. In the field, temperatures

during overwintering would alleviate dormancy (Fernández-Pascual et al., 2013b), probably slowly lowering T_b (Steadman and Pritchard, 2004, Pritchard et al., 1996, Pritchard et al., 1999, Batlla and Benech-Arnold, 2003, Porceddu et al., 2013) and reducing relative sensitivity to ΔT (Batlla et al., 2003, Benech-Arnold et al., 2000, Benech-Arnold et al., 1990a). Such anticipated changes require consideration. Finally, in situ soil temperatures can be expected to vary from those projected by air models, especially in fen soils where the buffer effect of the spring water is strongly felt (Fernández-Pascual et al., 2014). Overall, further complexity is needed in the development of cardinal-temperature models before they can encompass all the thermal determinants of seed germination and accurately predict plant regeneration under climate change.

ACKNOWLEDGEMENTS

T. García Gutiérrez, E. Lázaro González and P. Martín Pevida contributed to seed collection, which was authorized by the governments of Asturias and Castile-Leon. Special thanks to P. Lewis and O. Tierney for experimental support. J. Adams, N. Keogh and L. Queste also helped with laboratory work. We thank three anonymous reviewers for their improvements to the manuscript. The research was funded by the Department of Education and Science of the Government of Asturias (Convenio entre la Administración del Principado de Asturias y la Universidad de Oviedo, para el Desarrollo del Plan Estratégico de Viabilidad y Conversión a Campus de Excelencia Internacional) and had the institutional support of the Royal Botanic Gardens, Kew and Defra. E.F.P. was supported by the Government of Asturias (Grant BP09-107, Programa de Ayudas Predoctorales ‘Severo Ochoa’, Plan de Ciencia, Tecnología e Innovación del Principado de Asturias).

LITERATURE CITED

- Álvarez García MÁ, De Castro M, Cruz Guerrero R, Gómez Borrego V, Pérez Muñuzuri V, Stoll H. 2009. Clima. In: Anadón R, Roqueñí N, eds. CLIMAS. Evidencias y efectos potenciales del cambio climático en Asturias. Oviedo: Consejería de Medio Ambiente, Ordenación del Territorio e Infraestructuras.
- Baskin CC, Baskin JM. 1998. Seeds. Ecology, Biogeography, and Evolution of Dormancy and Germination. San Diego: Academic Press.
- Batlla D, Benech-Arnold RL. 2003. A quantitative analysis of dormancy loss dynamics in *Polygonum aviculare* L. seeds: development of a thermal time model based on changes in seed population thermal parameters. *Seed Science Research*, 13: 55-68.
- Batlla D, Benech-Arnold RL. 2004. A predictive model for dormancy loss in *Polygonum aviculare* L. seeds based on changes in population hydrotime parameters. *Seed Science Research*, 14: 277-286.
- Batlla D, Benech-Arnold RL. 2005. Changes in the light sensitivity of buried *Polygonum aviculare* seeds in relation to cold-induced dormancy loss: development of a predictive model. *New Phytologist*, 165: 445-452.
- Batlla D, Verges V, Benech-Arnold RL. 2003. A quantitative analysis of seed responses to cycle-doses of fluctuating temperatures in relation to dormancy: development of a thermal time model for

- Polygonum aviculare* L. seeds. *Seed Science Research*, 13: 197-207.
- Benech-Arnold RL, Ghersa CM, Sanchez RA, Insausti P. 1990a. A mathematical model to predict *Sorghum halepense* (L.) Pers. seedling emergence in relation to soil temperature. *Weed Research*, 30: 91-99.
- Benech-Arnold RL, Sánchez RA, Forcella F, Kruk BC, Ghersa CM. 2000. Environmental control of dormancy in weed seed banks in soil. *Field Crops Research*, 67: 105-122.
- Benech-Arnold RLG, C. M., Sanchez RA, Insausti P. 1990b. Temperature effects on dormancy release and germination rate in *Sorghum halepense* (L.) Pers. seeds: a quantitative analysis. *Weed Research*, 30: 81-89.
- Budelsky RA, Galatowitsch SM. 1999. Effects of moisture, temperature, and time on seed germination of five wetland *Carices*: implications for restoration. *Restoration Ecology*, 7: 86-97.
- Carta A, Bedini G, Müller J, Probert R. 2013. Comparative seed dormancy and germination of eight annual species of ephemeral wetland vegetation in a Mediterranean climate. *Plant Ecology*, 214: 339-349.
- Castroviejo S, Luceño M, Galán A, Jiménez Mejías P, Cabezas F, Medina L. 2007. *Flora iberica* vol. XVIII CYPERACEAE-PONTEDERIACEAE. Madrid: Real Jardín Botánico, CSIC.
- Cavieres LA, Arroyo MTK. 2000. Seed germination response to cold stratification period and thermal regime in *Phacelia secunda* (Hydrophyllaceae). *Plant Ecology*, 149: 1-8.
- Cochrane A, Daws MI, Hay F. 2011. Seed-based approach for identifying flora at risk from climate warming. *Austral Ecology*, 36: 923-935.
- Cochrane A, Yates CJ, Hoyle GL, Nicotra AB. 2014. Will among-population variation in seed traits improve the chance of species persistence under climate change? *Global Ecology and Biogeography*: n/a-n/a.
- Chiang GC, Bartsch M, Barua D, Nakabayashi K, Debieu M, Kronholm I, Koornneef M, Soppe WJ, Donohue K, De Meaux J. 2011. *DOG1* expression is predicted by the seed-maturation environment and contributes to geographical variation in germination in *Arabidopsis thaliana*. *Molecular Ecology*, 20: 3336-49.
- Daws MI, Lydall E, Chmielarz P, Leprince O, Pritchard HW, Matthews S, Thanos CA. 2004. Developmental heat sum influences recalcitrant seed traits in *Aesculus hippocastanum* across Europe. *New Phytologist*, 162: 157-166.
- Donohue K, de Casas RR, Burghardt L, Kovach K, Willis CG. 2010. Germination, postgermination adaptation, and species ecological ranges. In: Futuyma DJ, Shafer HB, Simberloff D, eds. *Annual Review of Ecology, Evolution, and Systematics* 41. Palo Alto: Annual Reviews.
- Donohue K, Dorn L, Griffith C, Kim E, Aguilera A, Polisetty CR, Schmitt J. 2005a. Environmental and genetic influences on the germination of *Arabidopsis thaliana* in the field. *Evolution*, 59: 740-757.
- Donohue K, Dorn L, Griffith C, Kim E, Aguilera A, Polisetty CR, Schmitt J. 2005b. The evolutionary ecology of seed germination of *Arabidopsis thaliana*: variable natural selection on germination timing. *Evolution*, 59: 758-770.
- Donohue K, Dorn L, Griffith C, Kim E, Aguilera A, Polisetty CR, Schmitt J. 2005c. Niche construction through germination cueing: life-history responses to timing of germination in *Arabidopsis thaliana*. *Evolution*, 59: 771-785.
- Fenner M. 1991. The effects of the parent environment on seed germinability. *Seed Science Research*, 1: 75-84.
- Fenner M, Thompson K. 2005. *The Ecology of Seeds*. Cambridge: Cambridge University Press.
- Fernández-Pascual E, Jiménez-Alfaro B. 2014. Phenotypic plasticity in seed germination correlates differentially with overwintering and flowering temperatures. *Seed Science Research*: in press.
- Fernández-Pascual E, Jiménez-Alfaro B, Caujapé-Castells J, Jaén-Molina R, Díaz TE. 2013a. A local dormancy cline is related to the seed maturation environment, population genetic composition and climate. *Annals of Botany*, 112: 937-945.
- Fernández-Pascual E, Jiménez-Alfaro B, Díaz TE. 2013b. The temperature dimension of the seed germination niche in fen wetlands. *Plant Ecology*, 214: 489-499.
- Fernández-Pascual E, Jiménez-Alfaro B, Hájek M, Díaz TE, Pritchard HW. 2014. The soil thermal buffer of fen wetlands and its implications for seed germination and climate resilience. *Folia Geobotanica et Phytotaxonomica*: in review.
- Figueroa R, Herms DA, Cardina J, Doohan D. 2010. Maternal environment effects on common groundsel (*Senecio vulgaris*) seed dormancy. *Weed Science*, 58: 160-166.
- Finch-Savage WE, Leubner-Metzger G. 2006. Seed dormancy and the control of germination. *New Phytologist*, 171: 501-523.
- Foley ME, Anderson JV, Chao WS, Doğramacı M, Horvath DP. 2010. Initial changes in the transcriptome of *Euphorbia esula* seeds induced to germinate with a combination of constant and diurnal alternating temperatures. *Plant Molecular Biology*, 73: 131-142.
- García-Huidobro J, Monteith JL, Squire GR. 1982. Time, temperature and germination of pearl millet (*Pennisetum typhoides* S. & H.): I. Constant temperature. *Journal of Experimental Botany*, 33: 288-296.
- Hardege SP. 2006. Predicting germination response to temperature. I. Cardinal-temperature models and subpopulation-specific regression. *Annals of Botany*, 97: 1115-1125.
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25: 1965-1978.
- Hoyle GL, Steadman KJ, Daws MI, Adkins SI. 2008. Pre- and post-harvest influences on seed dormancy status of an Australian Goodeniaceae species, *Goodenia fascicularis*. *Annals of Botany*, 102: 93-101.
- Hoyle GL, Venn SE, Steadman KJ, Good RB, McAuliffe EJ, Williams ER, Nicotra AB. 2013. Soil warming increases plant species richness but decreases germination from the alpine soil seed bank. *Global Change Biology*, 19: 1549-1561.
- Huarte HR. 2013. Mecanismos fisiológicos y moleculares involucrados en la terminación de la dormición de semillas expuestas a las temperaturas alternadas, PhD Thesis, University of Buenos Aires.
- Huarte HR, Benech-Arnold RL. 2005. Incubation under fluctuating temperatures reduces mean base water potential for seed germination in several non-cultivated species. *Seed Science Research*, 15: 89-97.
- Huarte HR, Benech-Arnold RL. 2010. Hormonal nature of seed responses to fluctuating temperatures in *Cynara cardunculus* (L.). *Seed Science Research*, 20: 39-45.
- IPCC. 2013. *Climate Change 2013. The Physical Science Basis. Summary for Policymakers*. Geneva: Intergovernmental Panel on Climate Change.
- Jones CD, Hughes JK, Bellouin N, Hardiman SC, Jones GS, Knight J, Liddicoat S, O'Connor FM, Andres RJ, Bell C, Boo KO, Bozzo A, Butchart N, Cadule P, Corbin KD, Doutriaux-Boucher M, Friedlingstein P, Gornall J, Gray L, Halloran PR, Hurtt G, Ingram WJ, Lamarque JF, Law RM, Meinshausen M, Osprey S, Palin EJ, Parsons Chini L, Raddatz T, Sanderson MG, Sellar AA, Schurer A, Valdes P, Wood N, Woodward S, Yoshioka M, Zerroukat M. 2011. The HadGEM2-ES implementation of CMIP5 centennial simulations. *Geoscientific Model Development*, 4: 543-570.
- Liu K, Baskin JM, Baskin CC, Bu H, Du G, Ma M. 2013. Effect of diurnal fluctuating versus constant temperatures on germination of 445 species from the eastern Tibet Plateau. *PloS one*, 8: e69364.
- Maas D. 1989. Germination characteristics of some plant species from calcareous fens in southern Germany and their implications for the seed bank. *Holarctic Ecology*, 12: 337-344.
- Molina A, Acedo C, Lence C, Alonso Felipe JI, Fernández A, Jiménez-Alfaro B, Robinson SG, Bueno Á, Llamas F. 2009. Estado de conservación y recatalogación UICN de *Carex diandra* Schrank en España. In: Llamas F, Acedo C, eds. *Botánica pirenaico-cantábrica en el siglo XXI*. León: University of León.
- Mondoni A, Rossi G, Orsenigo S, Probert RJ. 2012. Climate warming could shift the timing of seed germination in alpine plants. *Annals of Botany*, 110: 155-64.
- Morinaga T. 1926. Effect of alternating temperatures upon the germination of seeds. *American Journal of Botany*, 13: 141-158.
- Ooi MKJ. 2012. Seed bank persistence and climate change. *Seed Science Research*, 22: 53-60.

- Ooi MKJ, Auld TD, Denham AJ. 2009. Climate change and bet-hedging: interactions between increased soil temperatures and seed bank persistence. *Global Change Biology*, 15: 2375-2386.
- Orrù M, Mattana E, Pritchard HW, Bacchetta G. 2012. Thermal thresholds as predictors of seed dormancy release and germination timing: altitude-related risks from climate warming for the wild grapevine *Vitis vinifera* subsp. *sylvestris*. *Annals of Botany*, 110: 1651-1660.
- Patzelt A, Wild U, Pfadenhauer J. 2001. Restoration of wet fen meadows by topsoil removal: vegetation development and germination biology of fen species. *Restoration Ecology*, 9: 127-136.
- Porceddu M, Mattana E, Pritchard HW, Bacchetta G. 2013. Thermal niche for in situ seed germination by Mediterranean mountain streams: model prediction and validation for *Rhamnus persicifolia* seeds. *Annals of Botany*, 112: 1887-1897.
- Pritchard HW, Manger KR. 1990. Quantal response of fruit and seed germination rate in *Quercus robur* L. and *Castanea sativa* Mill, to constant temperatures and photon dose. *Journal of Experimental Botany*, 41: 1549-1557.
- Pritchard HW, Steadman KJ, Nash JV, Jones C. 1999. Kinetics of dormancy release and the high temperature germination response in *Aesculus hippocastanum* seeds. *Journal of Experimental Botany*, 50: 1507-1514.
- Pritchard HW, Tompsett PB, Manger KR. 1996. Development of a thermal time model for the quantification of dormancy loss in *Aesculus hippocastanum* seeds. *Seed Science Research*, 6: 127-135.
- Probert R, Smith R, Birch P. 1986. Germination responses to light and alternating temperatures in European populations of *Dactylis glomerata* L. V. The principle components of the alternating temperature requirement. *New Phytologist*, 102: 133-142.
- Probert RJ. 2000. The role of temperature in the regulation of seed dormancy and germination. In: Fenner M, ed. *Seeds: The Ecology of Regeneration in Plant Communities*. Wallingford: CABI.
- Qiu J, Bai Y, Coulman B, Romo JT. 2006. Using thermal time models to predict seedling emergence of orchardgrass (*Dactylis glomerata* L.) under alternating temperature regimes. *Seed Science Research*, 16: 261-271.
- Roberts EH, Totterdell S. 1981. Seed dormancy in *Rumex* species in response to environmental factors. *Plant, Cell & Environment*, 4: 97-106.
- Schütz W. 1997. Primary dormancy and annual dormancy cycles in seeds of six temperate wetland sedges. *Aquatic Botany*, 59: 75-85.
- Schütz W. 2000. Ecology of seed dormancy and germination in sedges (*Carex*). *Perspectives in Plant Ecology, Evolution and Systematics*, 3: 67-89.
- Schütz W, Rave G. 1999. The effect of cold stratification and light on the seed germination of temperate sedges (*Carex*) from various habitats and implications for regenerative strategies. *Plant Ecology*, 144: 215-230.
- Steadman KJ, Pritchard HW. 2004. Germination of *Aesculus hippocastanum* seeds following cold-induced dormancy loss can be described in relation to a temperature-dependent reduction in base temperature (T_b) and thermal time. *New Phytologist*, 161: 415-425.
- Thompson K, Grime JP. 1983. A comparative study of germination responses to diurnally-fluctuating temperatures. *Journal of Applied Ecology*, 20: 141-146.
- Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Webb DA. 1980. *Flora europaea* Volume 5. Cambridge: Cambridge University.
- Vasseur DA, DeLong JP, Gilbert B, Greig HS, Harley CDG, McCann KS, Savage V, Tunney TD, O'Connor MI. 2014. Increased temperature variation poses a greater risk to species than climate warming. *Proceedings of the Royal Society B: Biological Sciences*, 281.
- Vleeshouwers LM, Bouwmeester HJ, Karssen CM. 1995. Redefining seed dormancy: an attempt to integrate physiology and ecology. *Journal of Ecology*, 83: 1031-1037.
- Vuuren DP, Edmonds J, Kainuma M, Riahi K, Thomson A, Hibbard K, Hurtt GC, Kram T, Krey V, Lamarque J-F, Masui T, Meinshausen M, Nakicenovic N, Smith SJ, Rose SK. 2011. The representative concentration pathways: an overview. *Climatic Change*, 109: 5-31.
- Wagmann K, Hautekèete N-C, Piquot Y, Meunier C, Schmitt SE, Van Dijk H. 2012. Seed dormancy distribution: explanatory ecological factors. *Annals of Botany*, 110: 1205-1219.
- Walck JL, Hidayati SN, Dixon KW, Thompson K, Poschlod P. 2011. Climate change and plant regeneration from seed. *Global Change Biology*, 17: 2145-2161.
- Zhou Z, Bao W. 2014. Changes in seed dormancy of *Rosa multibracteata* Hemsl. & E. H. Wilson with increasing elevation in an arid valley in the eastern Tibetan Plateau. *Ecological Research*, 29: 693-700.

TABLE 1. Sources of variation in the thermal thresholds, base temperature for germination (T_b) and thermal time for 50% germination (θ_{50}). Results of a main effects Generalized Linear Model (gamma regression) that included the diurnal thermal amplitude (ΔT), population of origin and warmer phase (i.e., whether the warmer phase of the photoperiod was coupled to the light or the dark phase) as fixed predictors

Effect	T_b		θ_{50}	
	F	p	F	p
ΔT	36.626	<0.001	19.01	<0.001
Population	20.404	<0.001	1.943	0.172
Warmer phase	0.092	0.763	8.739	0.005




FIG. 1 Germination of *C. diandra* on the thermal gradient plate. Contour plots with points of equal percentage germination connected by germination isopleths. The shading represents 20 % germination increases, from lower (lighter filling) to higher (darker filling). Seeds were collected from a montane site (left panel) or subalpine site (right panel). The transversal axes of symmetry from the bottom-left corner to the top-right corner are constant thermal regimes. Points above and below the axes represent diurnally alternating temperatures regimes, with greatest amplitude at the top-left (warm night, cold day) and bottom-right (warm day, cold night) corners.




FIG. 2. Effect of the diurnal thermal amplitude (ΔT) in the base temperature for germination (T_b) and the thermal time for 50 % germination (θ_{50}) for seeds from montane and subalpine sites. The symbols represent the average values of T_b ($^{\circ}C$) and θ_{50} ($^{\circ}Cd$) at each ΔT regime; the lines are the logarithmic fits to the data.




FIG. 3. Simulated environmental temperature and base temperature (T_b) for germination at present and under the RCP2.6 and RCP8.5 scenarios. Each panel shows the results for one population under a given scenario. The solid line is the environmental monthly average temperature, the dotted line represents the T_b projected on the basis of the present diurnal thermal amplitude (ΔT) and the broken line is for the T_b projected in relation to the ΔT of the scenario.




FIG. 4. Annual accumulated environmental heat sum for germination ($EH\Sigma$) and thermal time required for 50 % germination (θ_{50}) at present (P) and under the RCP2.6 (2.6) and RCP8.5 (8.5) scenarios for montane and subalpine populations. The bars are the $EH\Sigma$ and the rectangles the θ_{50} . Left panels show the results of the simulation without taking into account the effect of the diurnal thermal oscillations (ΔT). Right panels show the results of the simulation considering future ΔT and its effect on the base temperature for germination and the θ_{50} .